

# P-stable Exponentially fitted Obrechkoff methods

Marnix Van Daele, G. Vanden Berghe

`Marnix.VanDaele@UGent.be`

Vakgroep Toegepaste Wiskunde en Informatica  
Universiteit Gent

# Outline

- Introduction on exponentially fitted (EF) methods
- 2-step Obrechkoff methods for  $y'' = f(x, y)$
- P-stable Obrechkoff methods for  $y'' = f(x, y)$
- P-stable EF Obrechkoff methods for  $y'' = f(x, y)$
- Conclusions

# Exponentially fitted methods

In the past 15 years, our research group has constructed modified versions of well-known

- linear multistep methods
- Runge-Kutta methods

Aim : build methods which perform very good when the solution has a known exponential or trigonometric behaviour.

# Linear multistep methods

A well known method to solve

$$y'' = f(y) \quad y(a) = y_a \quad y'(a) = y'_a$$

is the **Numerov method** (order 4)

$$y_{n+1} - 2y_n + y_{n-1} = \frac{h^2}{12} (f(y_{n-1}) + 10f(y_n) + f(y_{n+1}))$$

# Linear multistep methods

A well known method to solve

$$y'' = f(y) \quad y(a) = y_a \quad y'(a) = y'_a$$

is the **Numerov method** (order 4)

$$y_{n+1} - 2y_n + y_{n-1} = \frac{h^2}{12} (f(y_{n-1}) + 10f(y_n) + f(y_{n+1}))$$

Construction : impose  $\mathcal{L}[z(x); h] = 0$  for  $z(x) = 1, x, x^2, x^3, x^4$

where

$$\begin{aligned} \mathcal{L}[z(x); h] := & z(x+h) + \alpha_0 z(x) + \alpha_{-1} z(x-h) \\ & - h^2 (\beta_1 z''(x+h) + \beta_0 z''(x) + \beta_{-1} z''(x-h)) \end{aligned}$$

# Exponential fitting

Consider the initial value problem

$$y'' + \omega^2 y = g(y) \quad y(a) = y_a \quad y'(a) = y'_a.$$

If  $|g(y)| \ll |\omega^2 y|$  then

$$y(x) \approx \alpha \cos(\omega x + \phi)$$

To mimic this oscillatory behaviour, one could replace polynomials by trigonometric (in the complex case : exponential) functions.

# EF Numerov method

Construction : impose  $\mathcal{L}[z(x); h] = 0$  for

$$z(x) = 1, x, x^2, \sin(\omega x), \cos(\omega x)$$

$$\begin{aligned}\mathcal{L}[z(x); h] &:= z(x + h) + \alpha_0 z(x) + \alpha_{-1} z(x - h) \\ &\quad - h^2 (\beta_1 z''(x + h) + \beta_0 z''(x) + \beta_{-1} z''(x - h))\end{aligned}$$

# EF Numerov method

Construction : impose  $\mathcal{L}[z(x); h] = 0$  for

$$z(x) = 1, x, x^2, \sin(\omega x), \cos(\omega x)$$

$$\begin{aligned}\mathcal{L}[z(x); h] &:= z(x + h) + \alpha_0 z(x) + \alpha_{-1} z(x - h) \\ &\quad - h^2 (\beta_1 z''(x + h) + \beta_0 z''(x) + \beta_{-1} z''(x - h))\end{aligned}$$

$$y_{n+1} - 2y_n + y_{n-1} = h^2 (\lambda f(y_{n-1}) + (1 - 2\lambda) f(y_n) + \lambda f(y_{n+1}))$$

$$\lambda = \frac{1}{4 \sin^2 \frac{\theta}{2}} - \frac{1}{\theta^2} \qquad \qquad \theta := \omega h$$

# EF Numerov method

Construction : impose  $\mathcal{L}[z(x); h] = 0$  for

$$z(x) = 1, x, x^2, \sin(\omega x), \cos(\omega x)$$

$$\begin{aligned}\mathcal{L}[z(x); h] &:= z(x + h) + \alpha_0 z(x) + \alpha_{-1} z(x - h) \\ &\quad - h^2 (\beta_1 z''(x + h) + \beta_0 z''(x) + \beta_{-1} z''(x - h))\end{aligned}$$

$$y_{n+1} - 2y_n + y_{n-1} = h^2 (\lambda f(y_{n-1}) + (1 - 2\lambda) f(y_n) + \lambda f(y_{n+1}))$$

$$\lambda = \frac{1}{4 \sin^2 \frac{\theta}{2}} - \frac{1}{\theta^2} = \frac{1}{12} + \frac{1}{240} \theta^2 + \frac{1}{6048} \theta^4 + \dots \quad \theta := \omega h$$

# EF methods

Generalisation : to determine the coefficients of a method, we impose conditions on a linear functional. These conditions are related to

- polynomials :

$$\{x^q | q = 0, \dots, K\}$$

- exponential or trigonometric functions, multiplied with powers of  $x$  :

$$\{x^q \exp(\pm \mu x) | q = 0, \dots, P\}$$

or, with  $\omega = i \mu$ ,

$$\{x^q \cos(\omega x), x^q \sin(\omega x) | q = 0, \dots, P\}$$

# EF methods

Generalisation : to determine the coefficients of a method, we impose conditions on a linear functional. These conditions are related to

- polynomials :

$$\{x^q | q = 0, \dots, K\}$$

- exponential or trigonometric functions, multiplied with powers of  $x$  :

$$\{x^q \exp(\pm \mu x) | q = 0, \dots, P\}$$

or, with  $\omega = i \mu$ ,

$$\{x^q \cos(\omega x), x^q \sin(\omega x) | q = 0, \dots, P\}$$

Classical method :  $P = -1$

# Choice of $\omega$

- local optimization
  - based on local truncation error (lte)
  - $\omega$  is step-dependent
- global optimization
  - Preservation of geometric properties (periodicity, energy, ...)
  - $\omega$  is constant over the interval of integration

# Obrechkoff methods



Nikola Obrechkoff (1896-1963)

Obrechkoff methods (OM) : °1940 for quadrature

Milne : OM for solving diff. eq. : 1949

# Obrechkoff methods for $y'' = f(x, y)$

Two-step methods

$$y_{n+1} - 2y_n + y_{n-1} = \sum_{i=1}^m h^{2i} \left( \beta_{i0} y_{n+1}^{(2i)} + 2\beta_{i1} y_n^{(2i)} + \beta_{i0} y_{n-1}^{(2i)} \right)$$

# Obrechkoff methods for $y'' = f(x, y)$

Two-step methods

$$y_{n+1} - 2y_n + y_{n-1} = \sum_{i=1}^m h^{2i} \left( \beta_{i0} y_{n+1}^{(2i)} + 2\beta_{i1} y_n^{(2i)} + \beta_{i0} y_{n-1}^{(2i)} \right)$$

$$\mathcal{L}[z(x); h] := z(x+h) - 2z(x) + z(x-h)$$

$$- \sum_{i=1}^m h^{2i} \left( \beta_{i0} z^{(2i)}(x+h) + 2\beta_{i1} z^{(2i)}(x) + \beta_{i0} z^{(2i)}(x-h) \right)$$

symmetric method :  $\mathcal{L}[z(x); h] \equiv 0$  if  $z(x)$  is odd

# Obrechkoff methods for $y'' = f(x, y)$

Two-step methods

$$y_{n+1} - 2y_n + y_{n-1} = \sum_{i=1}^m h^{2i} \left( \beta_{i0} y_{n+1}^{(2i)} + 2\beta_{i1} y_n^{(2i)} + \beta_{i0} y_{n-1}^{(2i)} \right)$$

$$\mathcal{L}[z(x); h] := z(x+h) - 2z(x) + z(x-h)$$

$$- \sum_{i=1}^m h^{2i} \left( \beta_{i0} z^{(2i)}(x+h) + 2\beta_{i1} z^{(2i)}(x) + \beta_{i0} z^{(2i)}(x-h) \right)$$

symmetric method :  $\mathcal{L}[z(x); h] \equiv 0$  if  $z(x)$  is odd

$$\mathcal{L}[1; h] \equiv 0$$

# Obrechkoff methods for $y'' = f(x, y)$

Two-step methods

$$y_{n+1} - 2y_n + y_{n-1} = \sum_{i=1}^m h^{2i} \left( \beta_{i0} y_{n+1}^{(2i)} + 2\beta_{i1} y_n^{(2i)} + \beta_{i0} y_{n-1}^{(2i)} \right)$$

$$\mathcal{L}[z(x); h] := z(x+h) - 2z(x) + z(x-h)$$

$$- \sum_{i=1}^m h^{2i} \left( \beta_{i0} z^{(2i)}(x+h) + 2\beta_{i1} z^{(2i)}(x) + \beta_{i0} z^{(2i)}(x-h) \right)$$

symmetric method :  $\mathcal{L}[z(x); h] \equiv 0$  if  $z(x)$  is odd

$$\mathcal{L}[1; h] \equiv 0$$

order  $p \iff \mathcal{L}[x^q; h] = 0, q = 0, 1, \dots, p+1$

$$\text{lte} = C_{p+2} h^{p+2} y^{(p+2)}(x_n) + \mathcal{O}(h^{p+3}) \quad C_{p+2} = \frac{\mathcal{L}[x^{p+2}; h]}{(p+2)! h^{p+2}}$$

# Two-step OM

- $m = 1 : p = 4, C_6 = -\frac{1}{240}$  (Numerov method)

$$y_{n+1} - 2y_n + y_{n-1} = \frac{h^2}{12} \left( y_{n+1}^{(2)} + 10y_n^{(2)} + y_{n-1}^{(2)} \right)$$

# Two-step OM

- $m = 1 : p = 4, C_6 = -\frac{1}{240}$  (Numerov method)

$$y_{n+1} - 2y_n + y_{n-1} = \frac{h^2}{12} \left( y_{n+1}^{(2)} + 10y_n^{(2)} + y_{n-1}^{(2)} \right)$$

- $m = 2 : p = 8, C_{10} = \frac{59}{76204800}$

$$\begin{aligned} y_{n+1} - 2y_n + y_{n-1} &= \frac{h^2}{252} \left( 11y_{n+1}^{(2)} + 230y_n^{(2)} + 11y_{n-1}^{(2)} \right) \\ &\quad - \frac{h^4}{15120} \left( 13y_{n+1}^{(4)} - 626y_n^{(4)} + 13y_{n-1}^{(4)} \right) \end{aligned}$$

# Two-step OM

- $m = 3 : p = 12, C_{14} = -\frac{45469}{1697361329664000}$

$$\begin{aligned}y_{n+1} - 2y_n + y_{n-1} &= \\ \frac{h^2}{7788} \left( 229 y_{n+1}^{(2)} + 7330 y_n^{(2)} + 229 y_{n-1}^{(2)} \right) \\ - \frac{h^4}{25960} \left( 11 y_{n+1}^{(4)} - 1422 y_n^{(4)} + 11 y_{n-1}^{(4)} \right) \\ + \frac{h^6}{39251520} \left( 127 y_{n+1}^{(6)} + 4846 y_n^{(6)} + 127 y_{n-1}^{(6)} \right)\end{aligned}$$

- ...

method of order  $p = 4 m$

# Stability of two-step OM

$$y_{n+1} - 2y_n + y_{n-1} = \sum_{i=1}^m h^{2i} \left( \beta_{i0} y_{n+1}^{(2i)} + 2\beta_{i1} y_n^{(2i)} + \beta_{i0} y_{n-1}^{(2i)} \right)$$

applied to  $y'' = -\lambda^2 y$  gives

$$y_{n+1} - 2R_{mm}(\nu^2) y_n + y_{n-1} = 0 \quad \nu := \lambda h$$

$$R_{mm}(\nu^2) = \frac{1 + \sum_{i=1}^m (-1)^i \beta_{i1} \nu^{2i}}{1 + \sum_{i=1}^m (-1)^{i+1} \beta_{i0} \nu^{2i}}$$

# Stability of two-step OM

$$y_{n+1} - 2y_n + y_{n-1} = \sum_{i=1}^m h^{2i} \left( \beta_{i0} y_{n+1}^{(2i)} + 2\beta_{i1} y_n^{(2i)} + \beta_{i0} y_{n-1}^{(2i)} \right)$$

applied to  $y'' = -\lambda^2 y$  gives

$$y_{n+1} - 2R_{mm}(\nu^2) y_n + y_{n-1} = 0 \quad \nu := \lambda h$$

$$R_{mm}(\nu^2) = \frac{1 + \sum_{i=1}^m (-1)^i \beta_{i1} \nu^{2i}}{1 + \sum_{i=1}^m (-1)^{i+1} \beta_{i0} \nu^{2i}}$$

The stability function  $R_{mm}$  uniquely determines the method.

# Stability of two-step OM

$$y_{n+1} - 2y_n + y_{n-1} = \sum_{i=1}^m h^{2i} \left( \beta_{i0} y_{n+1}^{(2i)} + 2\beta_{i1} y_n^{(2i)} + \beta_{i0} y_{n-1}^{(2i)} \right)$$

applied to  $y'' = -\lambda^2 y$  gives

$$y_{n+1} - 2R_{mm}(\nu^2) y_n + y_{n-1} = 0 \quad \nu := \lambda h$$

$$R_{mm}(\nu^2) = \frac{1 + \sum_{i=1}^m (-1)^i \beta_{i1} \nu^{2i}}{1 + \sum_{i=1}^m (-1)^{i+1} \beta_{i0} \nu^{2i}}$$

The stability function  $R_{mm}$  uniquely determines the method.

A method has the interval of periodicity  $(0, \nu_0^2)$  if

$$|R_{mm}(\nu^2)| < 1 \text{ for } 0 < \nu^2 < \nu_0^2.$$

# Stability of two-step OM

$$y_{n+1} - 2y_n + y_{n-1} = \sum_{i=1}^m h^{2i} \left( \beta_{i0} y_{n+1}^{(2i)} + 2\beta_{i1} y_n^{(2i)} + \beta_{i0} y_{n-1}^{(2i)} \right)$$

applied to  $y'' = -\lambda^2 y$  gives

$$y_{n+1} - 2R_{mm}(\nu^2) y_n + y_{n-1} = 0 \quad \nu := \lambda h$$

$$R_{mm}(\nu^2) = \frac{1 + \sum_{i=1}^m (-1)^i \beta_{i1} \nu^{2i}}{1 + \sum_{i=1}^m (-1)^{i+1} \beta_{i0} \nu^{2i}}$$

The stability function  $R_{mm}$  uniquely determines the method.

A method has the interval of periodicity  $(0, \nu_0^2)$  if


$$|R_{mm}(\nu^2)| < 1 \text{ for } 0 < \nu^2 < \nu_0^2.$$

The method is **P-stable** if  $|R_{mm}(\nu^2)| \leq 1$  for all real  $\nu \neq 0$ .

# Stability of two-step OM

- $m = 1 : p = 4, C_6 = -\frac{1}{240}$  (Numerov)  $\nu_0^2 = 6$


$$y_{n+1} - 2y_n + y_{n-1} = \frac{h^2}{12} \left( y_{n+1}^{(2)} + 10y_n^{(2)} + y_{n-1}^{(2)} \right)$$



# Stability of two-step OM

- $m = 2 : p = 8, C_{10} = \frac{59}{76204800} \quad \nu_0^2 = 25.2$


$$\begin{aligned}y_{n+1} - 2y_n + y_{n-1} = \\ \frac{h^2}{252} \left( 11 y_{n+1}^{(2)} + 115 y_n^{(2)} + 11 y_{n-1}^{(2)} \right) \\ - \frac{h^4}{15120} \left( 13 y_{n+1}^{(4)} - 626 y_n^{(4)} + 13 y_{n-1}^{(4)} \right)\end{aligned}$$



# Stability of two-step OM

- $m = 3 : p = 12, C_{14} = -\frac{45469}{1697361329664000} \quad \nu_0^2 = 55.60\dots$

$$\begin{aligned}y_{n+1} - 2y_n + y_{n-1} &= \\&\quad \frac{h^2}{7788} \left( 229 y_{n+1}^{(2)} + 7330 y_n^{(2)} + 229 y_{n-1}^{(2)} \right) \\&\quad + \frac{h^4}{25960} \left( -11 y_{n+1}^{(4)} + 1422 y_n^{(4)} - 11 y_{n-1}^{(4)} \right) \\&\quad + \frac{h^6}{39251520} \left( 127 y_{n+1}^{(6)} + 4846 y_n^{(6)} + 127 y_{n-1}^{(6)} \right)\end{aligned}$$



# P-stable 2-step OM

Ananthakrishnaiah (1987)

idea : use some parameters to obtain P-stability.

# P-stable 2-step OM

Ananthakrishnaiah (1987)

idea : use some parameters to obtain P-stability.

$m = 3$  and  $m = 4$

# P-stable 2-step OM

Ananthakrishnaiah (1987)

idea : use some parameters to obtain P-stability.

$m = 3$  and  $m = 4$

e.g.  $m = 3$

$$\begin{aligned}y_{n+1} - 2y_n + y_{n-1} = \\ h^2 \left( \beta_{10} y_{n+1}^{(2)} + 2 \beta_{11} y_n^{(2)} + \beta_{10} y_{n-1}^{(2)} \right) \\ + h^4 \left( \beta_{20} y_{n+1}^{(4)} + 2 \beta_{21} y_n^{(4)} + \beta_{20} y_{n-1}^{(4)} \right) \\ + h^6 \left( \beta_{30} y_{n+1}^{(6)} + 2 \beta_{31} y_n^{(6)} + \beta_{30} y_{n-1}^{(6)} \right)\end{aligned}$$

# P-stable 2-step OM

Ananthakrishnaiah (1987)

idea : use some parameters to obtain P-stability.

$m = 3$  and  $m = 4$

e.g.  $m = 3$

$$\begin{aligned}y_{n+1} - 2y_n + y_{n-1} &= \\& h^2 \left( \beta_{10} y_{n+1}^{(2)} + 2 \beta_{11} y_n^{(2)} + \beta_{10} y_{n-1}^{(2)} \right) \\& + h^4 \left( \beta_{20} y_{n+1}^{(4)} + 2 \beta_{21} y_n^{(4)} + \beta_{20} y_{n-1}^{(4)} \right) \\& + h^6 \left( \beta_{30} y_{n+1}^{(6)} + 2 \beta_{31} y_n^{(6)} + \beta_{30} y_{n-1}^{(6)} \right)\end{aligned}$$

impose  $p = 6 : \{x^2, x^4, x^6\}$ ,

then 3 parameters ( $\beta_{20}$ ,  $\beta_{30}$  and  $\beta_{31}$ ) remain

# P-stable 2-step OM

Ananthakrishnaiah (1987)

idea : use some parameters to obtain P-stability.

$m = 3$  and  $m = 4$

e.g.  $m = 3$

$$\begin{aligned}y_{n+1} - 2y_n + y_{n-1} &= \\& h^2 \left( \beta_{10} y_{n+1}^{(2)} + 2 \beta_{11} y_n^{(2)} + \beta_{10} y_{n-1}^{(2)} \right) \\& + h^4 \left( \beta_{20} y_{n+1}^{(4)} + 2 \beta_{21} y_n^{(4)} + \beta_{20} y_{n-1}^{(4)} \right) \\& + h^6 \left( \beta_{30} y_{n+1}^{(6)} + 2 \beta_{31} y_n^{(6)} + \beta_{30} y_{n-1}^{(6)} \right)\end{aligned}$$

impose  $p = 6 : \{x^2, x^4, x^6\}$ ,

then 3 parameters ( $\beta_{20}$ ,  $\beta_{30}$  and  $\beta_{31}$ ) remain

$$R_{33} = \frac{1 - \beta_{11} \nu^2 + \beta_{21} \nu^4 - \beta_{31} \nu^6}{1 + \beta_{10} \nu^2 - \beta_{20} \nu^4 + \beta_{30} \nu^6}$$

let  $\beta_{31} = \beta_{30}$

# Ananthakrishnaiah's idea

Choose these 2 remaining parameters  $\beta_{20}, \beta_{30}$  such that the method becomes P-stable

$$|R_{33}| = \left| \frac{N_3}{D_3} \right| < 1$$

# Ananthakrishnaiah's idea


Choose these 2 remaining parameters  $\beta_{20}, \beta_{30}$  such that the method becomes P-stable

$$|R_{33}| = \left| \frac{N_3}{D_3} \right| < 1 \iff (D_3 - N_3)(D_3 + N_3) > 0$$

# Ananthakrishnaiah's idea

Choose these 2 remaining parameters  $\beta_{20}, \beta_{30}$  such that the method becomes P-stable

$$|R_{33}| = \left| \frac{N_3}{D_3} \right| < 1 \iff (D_3 - N_3)(D_3 + N_3) > 0$$



# Ananthakrishnaiah's idea

Find the solution for which the phase-lag,

$$\nu - \arccos R_{33} = \left( \frac{13}{604800} + \frac{13}{30} \beta_{30} + \frac{1}{40} \beta_{20} \right) \nu^7 + \mathcal{O}(\nu^9)$$


becomes minimal.

# Ananthakrishnaiah's idea

Find the solution for which the phase-lag,

$$\nu - \arccos R_{33} = \left( \frac{13}{604800} + \frac{13}{30} \beta_{30} + \frac{1}{40} \beta_{20} \right) \nu^7 + \mathcal{O}(\nu^9)$$

becomes minimal.




# Ananthakrishnaiah's idea

Find the solution for which the phase-lag,

$$\nu - \arccos R_{33} = \left( \frac{13}{604800} + \frac{13}{30} \beta_{30} + \frac{1}{40} \beta_{20} \right) \nu^7 + \mathcal{O}(\nu^9)$$

becomes minimal.



This gives  $(\beta_{30}, \beta_{20}) = \left( \frac{1}{14400}, -\frac{1}{600} \right)$ .


# Ananthakrishnaiah's idea

Ananthakrishnaiah  $m = 3 : p = 6$ , P-stable,  $C_8 = -\frac{1}{50400}$

$$\begin{aligned}y_{n+1} - 2y_n + y_{n-1} &= \\ \frac{h^2}{20} \left( y_{n+1}^{(2)} + 18y_n^{(2)} + y_{n-1}^{(2)} \right) \\ - \frac{h^4}{600} \left( y_{n+1}^{(4)} - 22y_n^{(4)} + y_{n-1}^{(4)} \right) \\ + \frac{h^6}{14400} \left( y_{n+1}^{(6)} + 2y_n^{(6)} + y_{n-1}^{(6)} \right)\end{aligned}$$

# Ananthakrishnaiah's idea

$$R_{33} = \frac{1 - \frac{9}{20} \nu^2 + \frac{11}{600} \nu^4 - \frac{1}{14400} \nu^6}{1 + \frac{1}{20} \nu^2 + \frac{1}{600} \nu^4 + \frac{1}{14400} \nu^6}$$



$$|R_{33}| = \left| \frac{N_3}{D_3} \right| < 1 \text{ since}$$

$$(D_3 - N_3)(D_3 + N_3) = \frac{\nu^2 (\nu^2 - 10)^2 (\nu^2 - 60)^2}{360000}$$

# P-stable 2-step OM

We were able to generalise Ananthakrishnaiah's idea in

M. Van Daele and G. Vanden Berghe, P-stable Obrechkoff methods of arbitrary order  
for second-order differential equations, Numerical Algorithms **44**, 2007, 115-131

Algorithm to construct a P-stable OM for a given  $m$  :

- impose order  $2m$
- $D_m - N_m$  and  $D_m + N_m$  should both be halves of perfect squares

# P-stable 2-step OM

We were able to generalise Ananthakrishnaiah's idea in

M. Van Daele and G. Vanden Berghe, P-stable Obrechkoff methods of arbitrary order  
for second-order differential equations, Numerical Algorithms **44**, 2007, 115-131

Algorithm to construct a P-stable OM for a given  $m$  :

- impose order  $2m$
- $D_m - N_m$  and  $D_m + N_m$  should both be halves of perfect squares

Theorem : the approximant  $R_{mm}(\nu^2)$ , obtained by generalising  
Ananthakrishnaiah's approach, is given by the real part of the  
 $(m, m)$ -Padé approximant of  $\exp(i\nu)$ .

# P-stable 2-step OM

We were able to generalise Ananthakrishnaiah's idea in

M. Van Daele and G. Vanden Berghe, P-stable Obrechkoff methods of arbitrary order  
for second-order differential equations, Numerical Algorithms **44**, 2007, 115-131

Algorithm to construct a P-stable OM for a given  $m$  :

- impose order  $2m$
- $D_m - N_m$  and  $D_m + N_m$  should both be halves of perfect squares

This leads to a system of **non-linear** equations.

Theorem : the approximant  $R_{mm}(\nu^2)$ , obtained by generalising  
Ananthakrishnaiah's approach, is given by the real part of the  
 $(m, m)$ -Padé approximant of  $\exp(i\nu)$ .

This leads to a system of **linear** equations.

# Example

$$\begin{aligned} R_{33}(\nu^2) &= \frac{1 - \frac{9}{20} \nu^2 + \frac{11}{600} \nu^4 - \frac{1}{14400} \nu^6}{1 + \frac{1}{20} \nu^2 + \frac{1}{600} \nu^4 + \frac{1}{14400} \nu^6} \\ &= \Re \left( \frac{1 + \frac{1}{2} i \nu - \frac{1}{10} \nu^2 - \frac{1}{120} i \nu^3}{1 - \frac{1}{2} i \nu - \frac{1}{10} \nu^2 + \frac{1}{120} i \nu^3} \right) \end{aligned}$$

where  $\frac{1 + \frac{1}{2} \nu + \frac{1}{10} \nu^2 + \frac{1}{120} \nu^3}{1 - \frac{1}{2} \nu + \frac{1}{10} \nu^2 - \frac{1}{120} \nu^3} = \exp(\nu) + \mathcal{O}(\nu^7)$

# Example

$$\begin{aligned} R_{33}(\nu^2) &= \frac{1 - \frac{9}{20} \nu^2 + \frac{11}{600} \nu^4 - \frac{1}{14400} \nu^6}{1 + \frac{1}{20} \nu^2 + \frac{1}{600} \nu^4 + \frac{1}{14400} \nu^6} \\ &= \Re \left( \frac{1 + \frac{1}{2} i \nu - \frac{1}{10} \nu^2 - \frac{1}{120} i \nu^3}{1 - \frac{1}{2} i \nu - \frac{1}{10} \nu^2 + \frac{1}{120} i \nu^3} \right) \\ &= \frac{\left(1 - \frac{1}{10} \nu^2\right)^2 - \left(\frac{1}{2} \nu - \frac{1}{120} \nu^3\right)^2}{\left(1 - \frac{1}{10} \nu^2\right)^2 + \left(\frac{1}{2} \nu - \frac{1}{120} \nu^3\right)^2} \end{aligned}$$

where  $\frac{1 + \frac{1}{2} \nu + \frac{1}{10} \nu^2 + \frac{1}{120} \nu^3}{1 - \frac{1}{2} \nu + \frac{1}{10} \nu^2 - \frac{1}{120} \nu^3} = \exp(\nu) + \mathcal{O}(\nu^7)$

$$|R_{33}(\nu^2)| = \left| \frac{\left(1 - \frac{1}{10} \nu^2\right)^2 - \left(\frac{1}{2} \nu - \frac{1}{120} \nu^3\right)^2}{\left(1 - \frac{1}{10} \nu^2\right)^2 + \left(\frac{1}{2} \nu - \frac{1}{120} \nu^3\right)^2} \right| \leq 1$$

# Conclusion

How does the P-stable Obrechkoff method

$$y_{n+1} - 2y_n + y_{n-1} = \sum_{i=1}^m h^{2i} \left( \beta_{i0} y_{n+1}^{(2i)} + 2\beta_{i1} y_n^{(2i)} + \beta_{i0} y_{n-1}^{(2i)} \right)$$

of order  $2m$  look like ?

$$\begin{cases} \beta_{i0} = (-1)^{i+1} a_i^2 + 2 \sum_{j=0}^{i-1} (-1)^{j+1} a_j a_{2i-j} & i = 1, \dots, m \\ \beta_{i1} = a_i^2 + 2 \sum_{j=0}^{i-1} a_j a_{2i-j} \end{cases}$$

$$\text{where } a_j = \begin{cases} \frac{\binom{m}{j}}{\binom{2m}{j}} & \text{for } 0 \leq j \leq m \\ 0 & \text{for } j > m \end{cases}$$

# P-stable Expon. fitted OM

How to obtain P-stable exponentially-fitted Obrechkoff methods ?

$$y_{n+1} - 2y_n + y_{n-1} = \sum_{i=1}^m h^{2i} \left( \beta_{i0} y_{n+1}^{(2i)} + 2\beta_{i1} y_n^{(2i)} + \beta_{i0} y_{n-1}^{(2i)} \right)$$

applied to  $y'' = -\lambda^2 y$  gives

$$y_{n+1} - 2R_{mm}(\theta, \nu^2) y_n + y_{n-1} = 0$$

with  $\theta := \omega h$  and  $\nu := \lambda h$

# P-stable Expon. fitted OM

How to obtain P-stable exponentially-fitted Obrechkoff methods ?

$$y_{n+1} - 2y_n + y_{n-1} = \sum_{i=1}^m h^{2i} \left( \beta_{i0} y_{n+1}^{(2i)} + 2\beta_{i1} y_n^{(2i)} + \beta_{i0} y_{n-1}^{(2i)} \right)$$

applied to  $y'' = -\lambda^2 y$  gives

$$y_{n+1} - 2R_{mm}(\theta, \nu^2) y_n + y_{n-1} = 0$$

with  $\theta := \omega h$  and  $\nu := \lambda h$

Padé-approximants  $\implies$  exponentially-fitted Padé approximants

# P-stable Expon. fitted OM

How to obtain P-stable exponentially-fitted Obrechkoff methods ?

$$y_{n+1} - 2y_n + y_{n-1} = \sum_{i=1}^m h^{2i} \left( \beta_{i0} y_{n+1}^{(2i)} + 2\beta_{i1} y_n^{(2i)} + \beta_{i0} y_{n-1}^{(2i)} \right)$$

applied to  $y'' = -\lambda^2 y$  gives

$$y_{n+1} - 2R_{mm}(\theta, \nu^2) y_n + y_{n-1} = 0$$

with  $\theta := \omega h$  and  $\nu := \lambda h$

Padé-approximants  $\implies$  exponentially-fitted Padé approximants

Construction of Polynomially-fitted  $(m, m)$  Padé approximants :

# P-stable Expon. fitted OM

How to obtain P-stable exponentially-fitted Obrechkoff methods ?

$$y_{n+1} - 2y_n + y_{n-1} = \sum_{i=1}^m h^{2i} \left( \beta_{i0} y_{n+1}^{(2i)} + 2\beta_{i1} y_n^{(2i)} + \beta_{i0} y_{n-1}^{(2i)} \right)$$

applied to  $y'' = -\lambda^2 y$  gives

$$y_{n+1} - 2R_{mm}(\theta, \nu^2) y_n + y_{n-1} = 0$$

with  $\theta := \omega h$  and  $\nu := \lambda h$

Padé-approximants  $\implies$  exponentially-fitted Padé approximants

Construction of Polynomially-fitted  $(m, m)$  Padé approximants :

$$\frac{P_m(x)}{P_m(-x)} = \exp(x) + \mathcal{O}(x^{2m+1})$$

# P-stable Expon. fitted OM

How to obtain P-stable exponentially-fitted Obrechkoff methods ?

$$y_{n+1} - 2y_n + y_{n-1} = \sum_{i=1}^m h^{2i} \left( \beta_{i0} y_{n+1}^{(2i)} + 2\beta_{i1} y_n^{(2i)} + \beta_{i0} y_{n-1}^{(2i)} \right)$$

applied to  $y'' = -\lambda^2 y$  gives

$$y_{n+1} - 2R_{mm}(\theta, \nu^2) y_n + y_{n-1} = 0$$

with  $\theta := \omega h$  and  $\nu := \lambda h$

Padé-approximants  $\implies$  exponentially-fitted Padé approximants

Construction of Polynomially-fitted  $(m, m)$  Padé approximants :

$$\frac{P_m(x)}{P_m(-x)} = \exp(x) + \mathcal{O}(x^{2m+1})$$

$$\iff \exp(x) P_m(-x) - P_m(x) = \mathcal{O}(x^{2m+1})$$

# P-stable Expon. fitted OM

How to obtain P-stable exponentially-fitted Obrechkoff methods ?

$$y_{n+1} - 2y_n + y_{n-1} = \sum_{i=1}^m h^{2i} \left( \beta_{i0} y_{n+1}^{(2i)} + 2\beta_{i1} y_n^{(2i)} + \beta_{i0} y_{n-1}^{(2i)} \right)$$

applied to  $y'' = -\lambda^2 y$  gives

$$y_{n+1} - 2R_{mm}(\theta, \nu^2) y_n + y_{n-1} = 0$$

with  $\theta := \omega h$  and  $\nu := \lambda h$

Padé-approximants  $\implies$  exponentially-fitted Padé approximants

Construction of Polynomially-fitted  $(m, m)$  Padé approximants :

$$\frac{P_m(x)}{P_m(-x)} = \exp(x) + \mathcal{O}(x^{2m+1})$$

$$\iff \exp(x) P_m(-x) - P_m(x) = \mathcal{O}(x^{2m+1})$$

$$\iff \frac{d^{2q}}{dx^{2q}} (\exp(x) P_m(-x) - P_m(x)) \Big|_{x=0} = 0 \quad q = 1, \dots, m$$

# EF Padé approximants

$$\mathcal{F}(x, t) = \exp(t x) V_m(-t x) - V_m(t x)$$

$$V_m(x) = 1 + \sum_{j=1}^m a_i x^i$$

$$\begin{cases} \frac{\partial^{2q}}{\partial x^{2q}} \mathcal{F}(x, t) \Big|_{(x,t)=(0,\theta)} = 0 & q = 1, \dots, K \\ \Re \left( \frac{\partial^q}{\partial t^q} \mathcal{F}(x, t) \Big|_{(x,t)=(i,\theta)} \right) = 0 & q = 0, \dots, P \end{cases}$$

where  $0 \leq K \leq m$  and  $P + K + 1 = m$ .

This leads to a system of  $m$  linear equations in the unknowns  $a_i$ ,

$$i = 1, \dots, m.$$

The EF  $(K, P)$  Padé approximant to  $\exp(\nu)$  is then given by

$$(K, P) \hat{P}_m^m(\nu) = V_m(\nu) / V_m(-\nu).$$

# EF Padé approximants : $m = 1$

$$\mathcal{F}(x, t) = \exp(t x) V_1(-t x) - V_1(t x) \quad V_m(x) = 1 + a_1 x$$

# EF Padé approximants : $m = 1$

$$\mathcal{F}(x, t) = \exp(t x) V_1(-t x) - V_1(t x) \quad V_m(x) = 1 + a_1 x$$

- $(K = 1, P = -1)$

$$\frac{\partial^2}{\partial x^2} \mathcal{F}(x, t) \Big|_{(x,t)=(0,\theta)} = 0 \iff \theta^2 (1 - 2 a_1) = 0 \iff a_1 = \frac{1}{2}$$

$${}^{(1,-1)}\hat{P}_1^1(\nu) = \frac{1 + \frac{1}{2} x}{1 - \frac{1}{2} x}$$

# EF Padé approximants : $m = 1$

$$\mathcal{F}(x, t) = \exp(t x) V_1(-t x) - V_1(t x) \quad V_m(x) = 1 + a_1 x$$

- $(K = 1, P = -1)$

$$\frac{\partial^2}{\partial x^2} \mathcal{F}(x, t) \Big|_{(x,t)=(0,\theta)} = 0 \iff \theta^2 (1 - 2 a_1) = 0 \iff a_1 = \frac{1}{2}$$

- $(K = 0, P = 0)$
- $$(1, -1) \hat{P}_1^1(\nu) = \frac{1 + \frac{1}{2} x}{1 - \frac{1}{2} x}$$

$$\begin{aligned} \Re(\mathcal{F}(i, \theta)) = 0 &\iff \Re \left( e^{i\theta} (1 - i a_1 \theta) - (1 + i a_1 \theta) \right) = 0 \\ &\iff a_1 = \frac{\sin \theta}{\theta (\cos \theta + 1)} = \frac{1}{2} \frac{\tan(\theta/2)}{\theta/2} \end{aligned}$$

$$(0, 0) \hat{P}_1^1(\nu) = \frac{1 + \frac{1}{2} \frac{\tan(\theta/2)}{\theta/2} x}{1 - \frac{1}{2} \frac{\tan(\theta/2)}{\theta/2} x}$$


# EF Padé approximants : $m = 1$

$m = 1$	
$(K, P)$	$a_1$
$(1, -1)$ $(0, 0)$	$\frac{1}{2}$ $\frac{\tan \frac{\theta}{2}}{\theta}$

# EF Padé approximants : $m = 1$

$m = 1$	
$(K, P)$	$a_1$
$(1, -1)$	$\frac{1}{2}$
$(0, 0)$	$\frac{1}{2} + \frac{\theta^2}{24} + \mathcal{O}(\theta^4)$

# EF Padé approximants : $m = 1$




# EF Padé approximants : $m = 2$

$m = 2$		
$(K, P)$	$a_1$	$a_2$
$(2, -1)$	$\frac{1}{2}$	$\frac{1}{12}$
$(1, 0)$	$\frac{1}{2}$	$\frac{2 \tan \frac{\theta}{2} - \theta}{2 \theta^2 \tan \frac{\theta}{2}}$
$(0, 1)$	$\frac{2 (1 - \cos \theta)}{(\theta + \sin \theta) \theta}$	$\frac{\theta - \sin \theta}{(\theta + \sin \theta) \theta^2}$


# EF Padé approximants : $m = 2$

$m = 2$		
$(K, P)$	$a_1$	$a_2$
$(2, -1)$	$\frac{1}{2}$	$\frac{1}{12}$
$(1, 0)$	$\frac{1}{2}$	$\frac{1}{12} + \frac{\theta^2}{720} + \mathcal{O}(\theta^4)$
$(0, 1)$	$\frac{1}{2} + \mathcal{O}(\theta^4)$	$\frac{1}{12} + \frac{\theta^2}{360} + \mathcal{O}(\theta^4)$

# EF Padé approximants : $m = 2$



$m = 3$



# The Stiefel-Bettis problem

$$z'' + z = 0.001 \exp(ix) \quad z(0) = 1 \quad z'(0) = 0.995i \quad 0 \leq x \leq 40\pi$$

equivalent real form

$$\begin{cases} u'' + u = 0.001 \cos(x), & u(0) = 1, \quad u'(0) = 0, \\ v'' + v = 0.001 \sin(x), & v(0) = 0, \quad v'(0) = 0.9995 . \end{cases}$$

$$z(x) = u(x) + i v(x)$$

$$u(x) = \cos(x) + 0.0005x \sin(x)$$


$$v(x) = \sin(x) - 0.0005x \cos(x)$$

$$\gamma(x) = \sqrt{u^2(x) + v^2(x)} = \sqrt{1 + (0.0005x)^2}$$

$$\gamma(40\pi) \quad h = 2^{-i}\pi \quad \omega = 1$$

# The Stiefel-Bettis problem


$$m = 1$$



$\circ : P = -1$        $\diamond : P = 0$

# The Stiefel-Bettis problem

$$m = 2$$




$\circ : P = -1$

$\diamond : P = 0$

$\square : P = 1$

# The Stiefel-Bettis problem

$$m = 3$$



$\circ : P = -1$

$\diamond : P = 0$

$\square : P = 1$

$\star : P = 2$

# Conclusion

two-step P-stable exponentially fitted Obrechkoff methods :

- for any given  $m$ , P-stable EF  $(K, P)$ -methods of order  $2m$  exist
- the construction is based on EF Padé approximants to the exponential function
- the coefficients depend on a parameter  $\theta$
- the coefficients are continuous functions of  $\theta$  iff  $P$  is odd
- numerical example was given to illustrate the theory