



Deferred correction based on exponentially-fitted mono-implicit Runge-Kutta methods

M. Van Daele

Department of Applied Mathematics, Computer Science and Statistics
Ghent University

SciCADE 2013
International Conference on Scientific Computation and
Differential Equations
September 16-20 2013

Outline

Exponential fitting

- Introduction

- EFRK methods

- EF as a correction technique

Deferred Correction

- What is deferred correction?

- The overall order of a DC-scheme

Mono-implicit Runge-Kutta methods

- Parameterized Runge-Kutta methods

- Mono-implicit Runge-Kutta methods

Deferred correction with EF

- Analysis of the error

- The scheme-level approach

- The method-level approach

- A priori knowledge



Exponential fitting

PART ONE

Exponential fitting

Aim :

to build methods which perform very good when the solution has a known exponential or trigonometric behaviour.

How : start from linear functional(s) and impose that for a set of functions \mathcal{S} the method produces exact results.

example : $\mathcal{S} = \{\cos \omega x, \sin \omega x, 1, x, x^2, \dots, x^{n-2}\}$

ω which is either real (trigonometric case) or purely imaginary (exponential case), **is determined from the expression for the local error.**

Example : Numerov method

$$y'' = f(y) \quad y(a) = y_a \quad y(b) = y_b$$

classical Numerov method :

$$y_{n+1} - 2y_n + y_{n-1} = \frac{1}{12} h^2 (f(y_{n+1}) + 10f(y_n) + f(y_{n-1}))$$

$$n = 1, 2, \dots, N \quad h = \frac{b-a}{N+1}$$

Construction :

impose $\mathcal{L}[z(x); h] = 0$ for $z(x) \in \mathcal{S} = \{1, x, x^2, x^3, x^4\}$ where

$$\begin{aligned} \mathcal{L}[z(x); h] := & z(x+h) + a_0 z(x) + a_{-1} z(x-h) \\ & - h^2 (b_1 z''(x+h) + b_0 z''(x) + b_{-1} z''(x-h)) \end{aligned}$$

$$\mathcal{L}[z(x); h] = -\frac{1}{240} h^6 z^{(6)}(x) + \mathcal{O}(h^8) \quad \implies \text{order 4}$$

Example : Numerov method

$$y'' = f(y) \quad y(a) = y_a \quad y(b) = y_b$$

classical Numerov method :

$$y_{n+1} - 2y_n + y_{n-1} = \frac{1}{12} h^2 (f(y_{n+1}) + 10f(y_n) + f(y_{n-1}))$$

$$n = 1, 2, \dots, N \quad h = \frac{b-a}{N+1}$$

Construction :

impose $\mathcal{L}[z(x); h] = 0$ for $z(x) \in \mathcal{S} = \{1, x, x^2, x^3, x^4\}$ where

$$\begin{aligned} \mathcal{L}[z(x); h] := & z(x+h) + a_0 z(x) + a_{-1} z(x-h) \\ & - h^2 (b_1 z''(x+h) + b_0 z''(x) + b_{-1} z''(x-h)) \end{aligned}$$

$$\mathcal{L}[z(x); h] = -\frac{1}{240} h^6 z^{(6)}(x) + \mathcal{O}(h^8) \quad \implies \text{order 4}$$

Example : Numerov method

$$y'' = f(y) \quad y(a) = y_a \quad y(b) = y_b$$

classical Numerov method :

$$y_{n+1} - 2y_n + y_{n-1} = \frac{1}{12} h^2 (f(y_{n+1}) + 10f(y_n) + f(y_{n-1}))$$

$$n = 1, 2, \dots, N \quad h = \frac{b-a}{N+1}$$

Construction :

impose $\mathcal{L}[z(x); h] = 0$ for $z(x) \in \mathcal{S} = \{1, x, x^2, x^3, x^4\}$ where

$$\begin{aligned} \mathcal{L}[z(x); h] := & z(x+h) + a_0 z(x) + a_{-1} z(x-h) \\ & - h^2 (b_1 z''(x+h) + b_0 z''(x) + b_{-1} z''(x-h)) \end{aligned}$$

$$\mathcal{L}[z(x); h] = -\frac{1}{240} h^6 z^{(6)}(x) + \mathcal{O}(h^8) \quad \implies \text{order 4}$$

Example : Numerov method

$$y'' = f(y) \quad y(a) = y_a \quad y(b) = y_b$$

classical Numerov method :

$$y_{n+1} - 2y_n + y_{n-1} = \frac{1}{12} h^2 (f(y_{n+1}) + 10f(y_n) + f(y_{n-1}))$$

$$n = 1, 2, \dots, N \quad h = \frac{b-a}{N+1}$$

Construction :

impose $\mathcal{L}[z(x); h] = 0$ for $z(x) \in \mathcal{S} = \{1, x, x^2, x^3, x^4\}$ where

$$\begin{aligned} \mathcal{L}[z(x); h] := & z(x+h) + a_0 z(x) + a_{-1} z(x-h) \\ & - h^2 (b_1 z''(x+h) + b_0 z''(x) + b_{-1} z''(x-h)) \end{aligned}$$

$$\mathcal{L}[z(x); h] = -\frac{1}{240} h^6 z^{(6)}(x) + \mathcal{O}(h^8) \quad \implies \text{order 4}$$

EF Numerov method

Construction : impose $\mathcal{L}[z(x); h] = 0$ for $z(x) \in \mathcal{S}$ with

$$\mathcal{S} = \{1, x, x^2, \sin(\omega x), \cos(\omega x)\}$$

$$\begin{aligned} \mathcal{L}[z(x); h] := & z(x+h) + a_0 z(x) + a_{-1} z(x-h) \\ & - h^2 (b_1 z''(x+h) + b_0 z''(x) + b_{-1} z''(x-h)) \end{aligned}$$

$$y_{n+1} - 2y_n + y_{n-1} = h^2 (\lambda f(y_{n-1}) + (1 - 2\lambda) f(y_n) + \lambda f(y_{n+1}))$$

$$\lambda = \frac{1}{4 \sin^2 \frac{\theta}{2}} - \frac{1}{\theta^2} = \frac{1}{12} + \frac{1}{240} \theta^2 + \frac{1}{6048} \theta^4 + \dots \quad \theta := \omega h$$

EF Numerov method

Construction : impose $\mathcal{L}[z(x); h] = 0$ for $z(x) \in \mathcal{S}$ with

$$\mathcal{S} = \{1, x, x^2, \sin(\omega x), \cos(\omega x)\}$$

$$\begin{aligned} \mathcal{L}[z(x); h] := & z(x+h) + a_0 z(x) + a_{-1} z(x-h) \\ & - h^2 (b_1 z''(x+h) + b_0 z''(x) + b_{-1} z''(x-h)) \end{aligned}$$

$$y_{n+1} - 2y_n + y_{n-1} = h^2 (\lambda f(y_{n-1}) + (1 - 2\lambda) f(y_n) + \lambda f(y_{n+1}))$$

$$\lambda = \frac{1}{4 \sin^2 \frac{\theta}{2}} - \frac{1}{\theta^2} = \frac{1}{12} + \frac{1}{240} \theta^2 + \frac{1}{6048} \theta^4 + \dots \quad \theta := \omega h$$

EF Numerov method

Construction : impose $\mathcal{L}[z(x); h] = 0$ for $z(x) \in \mathcal{S}$ with

$$\mathcal{S} = \{1, x, x^2, \sin(\omega x), \cos(\omega x)\}$$

$$\begin{aligned} \mathcal{L}[z(x); h] := & z(x+h) + a_0 z(x) + a_{-1} z(x-h) \\ & - h^2 (b_1 z''(x+h) + b_0 z''(x) + b_{-1} z''(x-h)) \end{aligned}$$

$$y_{n+1} - 2y_n + y_{n-1} = h^2 (\lambda f(y_{n-1}) + (1 - 2\lambda) f(y_n) + \lambda f(y_{n+1}))$$

$$\lambda = \frac{1}{4 \sin^2 \frac{\theta}{2}} - \frac{1}{\theta^2} = \frac{1}{12} + \frac{1}{240} \theta^2 + \frac{1}{6048} \theta^4 + \dots \quad \theta := \omega h$$

EF Numerov method

Construction : impose $\mathcal{L}[z(x); h] = 0$ for $z(x) \in \mathcal{S}$ with

$$\mathcal{S} = \{1, x, x^2, \sin(\omega x), \cos(\omega x)\}$$

or $\mathcal{S} = \{1, x, x^2, \exp(\mu x), \exp(-\mu x)\}$ $\mu := i\omega$

$$\begin{aligned} \mathcal{L}[z(x); h] := & z(x+h) + a_0 z(x) + a_{-1} z(x-h) \\ & - h^2 (b_1 z''(x+h) + b_0 z''(x) + b_{-1} z''(x-h)) \end{aligned}$$

$$y_{n+1} - 2y_n + y_{n-1} = h^2 (\lambda f(y_{n-1}) + (1 - 2\lambda) f(y_n) + \lambda f(y_{n+1}))$$

$$\lambda = \frac{1}{4 \sin^2 \frac{\theta}{2}} - \frac{1}{\theta^2} = \frac{1}{12} + \frac{1}{240} \theta^2 + \frac{1}{6048} \theta^4 + \dots \quad \theta := \omega h$$

EF Numerov method

Construction : impose $\mathcal{L}[z(x); h] = 0$ for $z(x) \in \mathcal{S}$ with

$$\mathcal{S} = \{1, x, x^2, \sin(\omega x), \cos(\omega x)\}$$

or $\mathcal{S} = \{1, x, x^2, \exp(\mu x), \exp(-\mu x)\}$ $\mu := i\omega$

$$\begin{aligned} \mathcal{L}[z(x); h] := & z(x+h) + a_0 z(x) + a_{-1} z(x-h) \\ & - h^2 (b_1 z''(x+h) + b_0 z''(x) + b_{-1} z''(x-h)) \end{aligned}$$

$$y_{n+1} - 2y_n + y_{n-1} = h^2 (\lambda f(y_{n-1}) + (1 - 2\lambda) f(y_n) + \lambda f(y_{n+1}))$$

$$\lambda = \frac{1}{4 \sin^2 \frac{\theta}{2}} - \frac{1}{\theta^2} = \frac{1}{12} + \frac{1}{240} \theta^2 + \frac{1}{6048} \theta^4 + \dots \quad \theta := \omega h$$

$$= -\frac{1}{4 \sinh^2 \frac{\nu}{2}} + \frac{1}{\nu^2} = \frac{1}{12} - \frac{1}{240} \nu^2 + \frac{1}{6048} \nu^4 + \dots \quad \nu := \mu h$$



Choice of ω

- local optimization
based on local truncation error (lte)
 ω is step-dependent
- global optimization
Preservation of geometric properties (periodicity, energy, ...)
 ω is constant over the interval of integration



Choice of ω

- local optimization
based on local truncation error (lte)
 ω is step-dependent
- global optimization
Preservation of geometric properties (periodicity, energy, ...)
 ω is constant over the interval of integration

EF Numerov method

$$\mathcal{S} = \{1, x, x^2, \sin(\omega x), \cos(\omega x)\}$$

$$y_{n+1} - 2y_n + y_{n-1} = h^2 (\lambda f(y_{n-1}) + (1 - 2\lambda) f(y_n) + \lambda f(y_{n+1}))$$

$$\mathcal{L}[z(x); h] = -\frac{1}{240} h^6 \left(z^{(6)}(x) + \omega^2 z^{(4)}(x) \right) + \mathcal{O}(h^8) \quad \implies \text{order 4}$$

A value for the parameter ω can be obtained from the expression for the lte :

$$y_n^{(6)}(x_n) + \omega_n^2 y_n^{(4)}(x_n) = 0.$$

Exponentially-fitted Runge-Kutta methods

For methods of collocation type (Gauss, LobattoIIIA, ...) **linear functionals** can be introduced

$$\left\{ \begin{array}{l} \mathcal{L}_i[y(x); h] = y(x + c_i h) - y(x) - h \sum_{j=1}^s a_{ij} y'(x + c_j h) \\ \qquad \qquad \qquad i = 1, \dots, s \\ \mathcal{L}[y(x); h] = y(x + h) - y(x) - h \sum_{i=1}^s b_i y'(x + c_i h). \end{array} \right.$$

A fitting space \mathcal{S} is introduced such that $\forall u \in \mathcal{S}$

$$\left\{ \begin{array}{l} \mathcal{L}_i[u(x); h] = 0 \quad i = 1, \dots, s \\ \mathcal{L}[u(x); h] = 0 \end{array} \right.$$

Trapezoidal rule

$$\mathcal{S} = \{1, x, x^2\}$$

$$\begin{array}{c|cc} 0 & 0 & 0 \\ 1 & \frac{1}{2} & \frac{1}{2} \\ \hline & \frac{1}{2} & \frac{1}{2} \end{array}$$

$$\text{lte}_{TR} = -\frac{h^3}{12}y^{(3)}(x_n) + \mathcal{O}(h^4)$$

Exponentially-fitted trapezoidal rule

$$\mathcal{S} = \{1, \exp(\mu x), \exp(-\mu x)\}$$

0	0	0
1	$\frac{s(\mu/2)}{c(\mu/2) \mu h}$	$\frac{s(\mu/2)}{c(\mu/2) \mu h}$
	$\frac{s(\mu/2)}{c(\mu/2) \mu h}$	$\frac{s(\mu/2)}{c(\mu/2) \mu h}$

$$c(t) := \cosh(th)$$

$$s(t) := \sinh(th)$$

$$\text{lte}_{EFTR} = -\frac{h^3}{12} \left(y^{(3)}(x_n) - \mu^2 y'(x_n) \right) + \mathcal{O}(h^4)$$



EF as a correction technique

In the past, EF has been successfully applied as a **correction technique**, see e.g.

- G. VANDEN BERGHE, H. DE MEYER, ON A CORRECTION OF NUMEROV-LIKE EIGENVALUE APPROXIMATIONS FOR STURM-LIOUVILLE PROBLEMS, JCAM **37** (1991) 179-186.

EF as a correction technique ...

- Sturm-Liouville problem in normal form :

$$-y'' + q(x)y = \lambda y \quad y(0) = y(\pi) = 0 \quad (1)$$

- discretisation (Numerov method) :

$$A v + B Q v = \Sigma B v \quad (2)$$

of size N , whereby $(N + 1)h = \pi$

- The eigenvalues $\lambda_1 < \lambda_2 < \lambda_3 < \dots$ of (1) are approximated by the eigenvalues $\Lambda_1 < \Lambda_2 < \Lambda_3 < \dots < \Lambda_N$ of (2)
- $\|\Lambda_k - \lambda_k\| = \mathcal{O}(k^6 h^4)$
- with EF, the error can be reduced seriously ...



... EF as a correction technique

To obtain a better approximation for λ_k

- compute approximation y_k of the exact eigenvector v_k
- compute ω_n^2 from the error-expression of the EF Numerov method $y_k^{(6)}(t_n) + \omega_n^2 y_k^{(4)}(t_n) = 0, n = 1, \dots, N$
- discretisation (EF Numerov method) :

$$A_{EF} v + B_{EF} Q v_{EF} = \Sigma_{EF} B_{EF} v_{EF} \quad (3)$$

- the eigenvalues $\lambda_1 < \lambda_2 < \lambda_3 < \dots$ of (1) are approximated by the eigenvalues $\Lambda_{EF,1} < \Lambda_{EF,2} < \Lambda_{EF,3} < \dots < \Lambda_{EF,N}$ of (3)
- $\|\Lambda_{EF,k} - \lambda_k\| = \mathcal{O}(k^3 h^4)$



Deferred Correction

PART TWO

Deferred correction (FOX (1947), LINDBERG (1980))

Consider the nonlinear two-point boundary value problem

$$\frac{dy}{dx} = f(x, y(x)), \quad g(y(a), y(b)) = 0, \quad a \leq x \leq b$$

- (1) discretisation (with Runge-Kutta method ϕ_p of order p) leads to a system of nonlinear algebraic conditions

$$\phi_p(\eta) = 0$$

- (2) Use second Runge-Kutta formula ϕ_{p+r} of order $p+r$ as follows

$$\phi_p(\bar{\eta}) = -\phi_{p+r}(\eta)$$

$-\phi_{p+r}(\eta)$ provides an estimate of the local truncation error of the lower order formula ϕ_p .

Iterated deferred correction

- This idea can be extended to allow for more deferred correction iterations.
- In the widely used code `TWPBVP.f` (Cash, Mazzia), the algorithm is

$$\phi_4(\eta) = 0$$

$$\phi_4(\bar{\eta}) = -\phi_6(\eta)$$

$$\phi_4(\bar{\bar{\eta}}) = -\phi_6(\eta) - \phi_8(\bar{\eta})$$

where the Runge-Kutta formulae ϕ_4 , ϕ_6 and ϕ_8 are MIRK formulae of order 4, 6 and 8, respectively.



Iterated deferred correction

- Computational advantages:
 - the solutions $\eta, \bar{\eta}, \dots$ are computed on the same grid, and this generally leads to a **large saving in the cost of the linear algebra**
 - The **local error estimates** $\bar{\eta} - \eta, \bar{\bar{\eta}} - \bar{\eta}$ are immediately available at **no extra cost**
- If the error is less than a user supplied tolerance, then the most accurate solution is accepted. If the error criterion is not satisfied, then the **mesh is refined** and the deferred correction scheme is applied on the new mesh.

The overall order of a DC-scheme

The criteria under which the numerical solution of a deferred correction scheme reaches a certain order, were established by Skeel for the problem

$$\phi(\eta) = 0 \tag{4}$$

$$\phi(\bar{\eta}) = \psi(\eta). \tag{5}$$

ϕ : basic discretization scheme $\psi(\eta)$: estimate of the residual

We assume that the solutions η and $\bar{\eta}$ are computed on a grid

$$\pi : \mathbf{a} = \mathbf{x}_1 < \mathbf{x}_2 < \dots < \mathbf{x}_{N+1} = \mathbf{b}.$$

Δy : restriction of continuous solution $y(x)$ to the finite grid π

The overall order of a DC-scheme

SKEEL, R. D., A THEORETICAL FRAMEWORK FOR PROVING ACCURACY RESULTS FOR DEFERRED CORRECTIONS, SINUM **19** (1982) 171-196

Let ϕ be a stable numerical method and assume that the following conditions hold for the deferred correction scheme (4), (5) :

- (i) $\|\eta - \Delta y\| = O(h^p)$
- (ii) $\|\psi(\Delta y) - \phi(\Delta y)\| = O(h^{r+p})$
- (iii) $\psi(\Delta w) = O(h^r)$

for arbitrary functions w having at least r continuous derivatives and $\|\cdot\|$ is a suitable finite norm. If $\phi(\bar{\eta}) = \psi(\eta)$ then

$$\|\bar{\eta} - \Delta y\| = O(h^{r+p}).$$



The overall order of a DC-scheme

The **real difficulty** in satisfying the conditions of Skeel's theorem comes from **condition (iii)**.

- in general $r = 1$
- to obtain $r = 2$ ψ has to be written in a special, parameterised form



Mono-implicit Runge-Kutta methods

PART THREE

Parameterized Runge-Kutta methods

Any Runge-Kutta method

$$y_{k+1} = y_k + h \sum_{i=1}^s b_i f(x_k + c_i h, Y_i)$$

$$Y_i = y_k + h \sum_{j=1}^s a_{ij} f(x_k + c_j h, Y_j), \quad i = 1 \dots s$$

can be written in a **parameterized form**

$$y_{k+1} = y_k + h \sum_{i=1}^s b_i f(x_k + c_i h, Y_i)$$

$$Y_i = (1 - v_i) y_k + v_i y_{k+1} + h \sum_{j=1}^s x_{ij} f(x_k + c_j h, Y_j), \quad i = 1 \dots s$$

Parameterized Runge-Kutta methods

Any Runge-Kutta method

$$\begin{array}{c|c} c & A \\ \hline & b^T \end{array}$$

can be written in a parameterized form

$$\begin{array}{c|c|c} c & v & X \\ \hline & & b^T \end{array}$$

whereby

$$X = A - v b^T$$

Example 1: the trapezoidal rule

$$y_{k+1} = y_k + \frac{h}{2} (f(x_k, y_k) + f(x_{k+1}, y_{k+1}))$$

- written as a Runge-Kutta method (= 2-stage Lobatto IIIA)

$$\begin{array}{l}
 y_{k+1} = y_k + \frac{h}{2} (f(x_k, Y_1) + f(x_{k+1}, Y_2)) \\
 Y_1 = y_k \\
 Y_2 = y_k + \frac{h}{2} (f(x_k, Y_1) + f(x_{k+1}, Y_2))
 \end{array}
 \quad
 \begin{array}{c|cc}
 0 & 0 & 0 \\
 1 & \frac{1}{2} & \frac{1}{2} \\
 \hline
 & \frac{1}{2} & \frac{1}{2}
 \end{array}$$

- written as a PIRK method

$$\begin{array}{l}
 y_{k+1} = y_k + \frac{h}{2} (f(x_k, Y_1) + f(x_{k+1}, Y_2)) \\
 Y_1 = y_k \\
 Y_2 = y_{k+1}
 \end{array}
 \quad
 \begin{array}{c|cc|cc}
 0 & 0 & 0 & 0 \\
 1 & 1 & 0 & 0 \\
 \hline
 & & \frac{1}{2} & \frac{1}{2}
 \end{array}$$

Example 2: the 3-stage Lobatto IIIA method

0	0	0	0
$\frac{1}{2}$	$\frac{5}{24}$	$\frac{1}{3}$	$-\frac{1}{24}$
1	$\frac{1}{6}$	$\frac{2}{3}$	$\frac{1}{6}$
	$\frac{1}{6}$	$\frac{2}{3}$	$\frac{1}{6}$

- PIRK version 1:

0	0	0	0	0
$\frac{1}{2}$	0	$\frac{5}{24}$	$\frac{1}{3}$	$-\frac{1}{24}$
1	1	0	0	0
		$\frac{1}{6}$	$\frac{4}{6}$	$\frac{1}{6}$

$$y_{k+1} = y_k + \frac{h}{6} (f(x_k, y_k) + 4f(x_k + \frac{h}{2}, Y_2) + f(x_{k+1}, y_{k+1}))$$

$$Y_2 = y_k + \frac{h}{24} (5f(x_k, y_k) + 8f(x_k + \frac{h}{2}, Y_2) - f(x_{k+1}, y_{k+1}))$$

Example 2: the 3-stage Lobatto IIIA method

0	0	0	0
$\frac{1}{2}$	$\frac{5}{24}$	$\frac{1}{3}$	$-\frac{1}{24}$
1	$\frac{1}{6}$	$\frac{2}{3}$	$\frac{1}{6}$
	$\frac{1}{6}$	$\frac{2}{3}$	$\frac{1}{6}$

- PIRK version 2:

0	0	0	0	0
1	1	0	0	0
$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{8}$	$-\frac{1}{8}$	0
		$\frac{1}{6}$	$\frac{1}{6}$	$\frac{4}{6}$

$$y_{k+1} = y_k + \frac{h}{6} (f(x_k, y_k) + 4f(x_k + \frac{h}{2}, Y_2) + f(x_{k+1}, y_{k+1}))$$

$$Y_2 = \frac{1}{2}y_k + \frac{1}{2}y_{k+1} + \frac{h}{8} (f(x_k, y_k) - f(x_{k+1}, y_{k+1}))$$

Example 2: the 3-stage Lobatto IIIA method

0	0	0	0
$\frac{1}{2}$	$\frac{5}{24}$	$\frac{1}{3}$	$-\frac{1}{24}$
1	$\frac{1}{6}$	$\frac{2}{3}$	$\frac{1}{6}$
	$\frac{1}{6}$	$\frac{2}{3}$	$\frac{1}{6}$

- PIRK version 2:

0	0	0	0	0
1	1	0	0	0
$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{8}$	$-\frac{1}{8}$	0
		$\frac{1}{6}$	$\frac{1}{6}$	$\frac{4}{6}$

MIRK: mono-implicit RK

$$y_{k+1} = y_k + \frac{h}{6} (f(x_k, y_k) + 4f(x_k + \frac{h}{2}, Y_2) + f(x_{k+1}, y_{k+1}))$$

$$Y_2 = \frac{1}{2}y_k + \frac{1}{2}y_{k+1} + \frac{h}{8} (f(x_k, y_k) - f(x_{k+1}, y_{k+1}))$$



Parameterized Runge-Kutta methods

- For a given Runge-Kutta method, there thus exist **infinitely many PIRK versions**.
- As **standalone method**, all parameterized versions of a given Runge-Kutta method give (in exact arithmetic) the **same** result
- But in a **deferred correction** setting, all PIRK versions are quite **different**.

Example : $y' = y \quad y(0) = 1 \quad y(x) = \exp(x)$

- ϕ : trapezoidal rule: $-y_1 + y_0 + \frac{h}{2}(y_0 + y_1) = 0$

$$\implies y_1 = \frac{1 + \frac{h}{2}}{1 - \frac{h}{2}} = \exp(h) + \frac{1}{12}h^3 + \mathcal{O}(h^4)$$

- ψ : PIRK version 1 of Lobatto IIIA

$$-\bar{y}_1 + \bar{y}_0 + \frac{h}{2}(\bar{y}_0 + \bar{y}_1) = -\text{est}_1$$

$$\text{est}_1 = -y_1 + y_0 + \frac{h}{6}(y_0 + 4Y_2 + y_1)$$

$$Y_2 = y_0 + \frac{h}{24}(5y_0 + 8Y_2 - y_1)$$

$$\implies \bar{y}_1 = \exp(h) - \frac{1}{36}h^4 + \mathcal{O}(h^5)$$

Example : $y' = y$ $y(0) = 1$ $y(x) = \exp(x)$

- ϕ : trapezoidal rule: $-y_1 + y_0 + \frac{h}{2}(y_0 + y_1) = 0$

$$\implies y_1 = \frac{1 + \frac{h}{2}}{1 - \frac{h}{2}} = \exp(h) + \frac{1}{12}h^3 + \mathcal{O}(h^4)$$

- ψ : PIRK version 2 of Lobatto IIIA

$$-\bar{y}_1 + \bar{y}_0 + \frac{h}{2}(\bar{y}_0 + \bar{y}_1) = -\text{est}_2$$

$$\text{est}_2 = -y_1 + y_0 + \frac{h}{6}(y_0 + 4Y_2 + y_1)$$

$$Y_2 = (y_0 + y_1)/2$$


$$\implies \bar{y}_1 = \exp(h) - \frac{1}{120}h^5 + \mathcal{O}(h^6)$$

Mono-implicit Runge-Kutta methods

Restricting X to **lower triangular** matrices only reveals the class of mono-implicit Runge-Kutta (MIRK) methods.

A MIRK method is only implicit in the next knot point, which makes such a method very suitable for solving boundary value problems and using deferred correction, while maintaining a good stability:

- if ϕ_p is an s -stage MIRK method, then the **dimension of the system** to be solved in each step of the deferred correction process is equal to the **number of gridpoints** in the integration interval, whereas for a general PIRK method the dimension will be s times larger
- if ψ is a MIRK method, then the error estimator can be computed **explicitly**, i.e. at a very low cost.



Deferred correction with EF

PART FOUR

joint work with Davy Hollevoet

Exponentially-fitted mono-implicit trapezoidal rule

$$\mathcal{S} = \{1, \exp(\omega x), \exp(-\omega x)\}$$

0	0	0	0
1	1	0	0
		$\frac{s(\omega/2)}{c(\omega/2)\omega h}$	$\frac{s(\omega/2)}{c(\omega/2)\omega h}$

$$c(t) := \cosh(th)$$

$$s(t) := \sinh(th)$$

$$\text{lte}_{EFTR} = -\frac{h^3}{12} \left(y^{(3)}(x_k) - \omega^2 y'(x_k) \right) + \mathcal{O}(h^4)$$

Exponentially-fitted mono-implicit Lobatto IIIA method


$$\mathcal{S} = \{1, x, \exp(\mu x), \exp(-\mu x)\}$$

0	0	0	0	0
1	1	0	0	0
$\frac{1}{2}$	$\frac{1}{2}$	$\frac{s(\mu/2)}{2\mu h(1+c(\mu/2))}$	$\frac{s(\mu/2)}{-2\mu h(1+c(\mu/2))}$	0
		$\frac{\mu h - 2s(\mu/2)}{2\mu h(1-c(\mu/2))}$	$\frac{\mu h - 2s(\mu/2)}{2\mu h(1-c(\mu/2))}$	$\frac{2s(\mu/2) - c(\mu/2)\mu h}{\mu h(1-c(\mu/2))}$

$$c(t) := \cosh(th)$$

$$s(t) := \sinh(th)$$

$$\text{lte}_{EFLob} = -\frac{h^5}{2880} \left(y^{(5)}(x_k) - \mu^2 y^{(3)}(x_k) - 5f_y \left(y^{(4)}(x_k) - \mu^2 y^{(2)}(x_k) \right) \right) + \mathcal{O}(h^6)$$




Analysis of the error

As it is the spirit of EF methods, we want to increase the accuracy of the overall method by the introduction of EF methods

This requires a detailed analysis of the expression for the error of a DC scheme.

Analysis in terms of [P-series](#) (details can be found in Ph.D. thesis of Davy Hollevoet)



Residual of a RK method

$\rho(t)$	ϕ^1
1	0
\vdots	\vdots
p	0
$p+1$	$\neq 0$

Example: trapezoidal rule

$\rho(t)$	t	$\phi^1(t)$
1	•	0
2	↓, ↑	0, 0
3	↙, ↘, ↙, ↓, ↑	$-\frac{1}{2'}, -\frac{1}{2'}, -\frac{1}{2'}, -\frac{1}{2'}, -\frac{1}{2}$

$$\begin{aligned}
 \phi(y(x_k), y(x_{k+1})) &= -\frac{1}{2} \frac{h^3}{6} \left[F(\text{↙}) + 2F(\text{↘}) + F(\text{↙}) \right. \\
 &\quad \left. + F(\text{↓}) + F(\text{↑}) \right] (x_k; y(x_k)) + \mathcal{O}(h^4) \\
 &= -\frac{1}{12} h^3 y^{(3)}(x_k) + \mathcal{O}(h^4).
 \end{aligned}$$

Residual of an EF RK method

$\rho(t)$	$\phi[\omega]^1(t)$	$1 \times$	$+\omega_h \times$	\dots	$+\omega_h^p \times$
1	$\mathcal{O}(\omega_h^p)$	0	0	\dots	$\phi_{t,p}^1$
\vdots	\vdots	\vdots	\vdots	\ddots	
p	$\mathcal{O}(\omega_h^1)$	0	$\phi_{t,1}^1$		
$p+1$	$\mathcal{O}(1)$	$\phi_{t,0}^1$			

$$\begin{aligned}
 \rho(y(x_k), y(x_{k+1})) &= B^*(\phi[\omega]^1, x_k, y(x_k)) \\
 &= \sum_{t \in T_B} \frac{h^{\rho(t)}}{\rho(t)!} \alpha(t) \phi[\omega]^1(t) F(t)(x_k; y(x_k))
 \end{aligned}$$

The residual must be $\mathcal{O}(h^{p+1})$, which implies that $\phi[\omega_h]^1(t)$ must be $\mathcal{O}(h^{p+1-\rho(t)})$ for all trees of order $\leq p+1$. The functions under consideration actually depend on $\omega_h := h\omega$, which leads us to the bounds in [Table 4.3](#).

Example: The EF trapezoidal rule

$\rho(t)$	t	$\phi[\omega]^1$		
		$1 \times$	$+\omega h \times$	$+\omega^2 h^2 \times$
1	•	0	0	$\frac{1}{12}$
2	⋮, ⋮	0, 0	0, 0	
3	∇, ∇, ∇, ⋮, ⋮	$-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}$		

We find^a that the leading residual term is provided by

$$b_3(\phi[\omega]^1) = \omega^2 \frac{1}{12} F(\cdot) - \frac{1}{2} \frac{1}{3!} \left[F(\nabla) + 2F(\nabla) + F(\nabla) + F(\ddot{\cdot}) + F(\ddot{\cdot}) \right].$$

If at every knot point x_k an appropriate value for ω can be found such that

$$b_3(\phi[\omega]^1)(x_k; y(x_k)) = 0, \quad (4.24)$$


then those values can be used to increase the accuracy of the exponentially fitted trapezoid rule to order 3 for the problem at hand. This is due to the fact that the right-hand side of (4.24) is the coefficient of h^3 in a series expansion of the residual, perhaps more recognisable in terms of total derivatives:

$$b_3(\phi[\omega]^1)(x_k; y(x_k)) = -\frac{1}{12} y^{(3)}(x_k) + \omega^2 \frac{1}{12} y'(x_k).$$




Residual of a DC scheme

$\rho(t)$	ϕ^1	$-\psi^v$
1	0	0
\vdots	\vdots	\vdots
p	0	0
$p+1$	ϕ_t^1	ϕ_t^1
\vdots	\vdots	\vdots
q	ϕ_t^1	ϕ_t^1
$q+1$	ϕ_t^1	$\neq \phi_t^1$



Residual of an EF DC scheme

$\rho(t)$	$\phi[\omega]^1$				$-\psi[\mu]^v[\omega]$			
	$1 \times$	$+h \times$	\dots	$+h^p \times$	$1 \times$	$+h \times$	\dots	$+h^p \times$
1	0	0	\dots	$\omega^p \phi_{t,p}^1$	0	0	\dots	$\omega^p \phi_{t,p}^1$
\vdots	\vdots	\vdots	\ddots	\vdots	\vdots	\vdots	\ddots	\vdots
p	0	$\omega \phi_{t,1}^1$	\dots	$\omega^p \phi_{t,p}^1$	0	$\omega \phi_{t,1}^1$	\dots	$\omega^p \phi_{t,p}^1$
$p+1$	$\phi_{t,0}^1$	$\omega \phi_{t,1}^1$	\dots	$\omega^p \phi_{t,p}^1$	$\phi_{t,0}^1$	$\omega \phi_{t,1}^1$	\dots	$\psi_{t,p}^v$
\vdots	\vdots	\vdots	\ddots		\vdots	\vdots	\ddots	
q	$\phi_{t,0}^1$	$\omega \phi_{t,1}^1$			$\phi_{t,0}^1$	$\psi_{t,1}^v$		
$q+1$	$\phi_{t,0}^1$				$\psi_{t,0}^v$			



The overall error of TR - Radaul

$\rho(t)$	t	$\phi[\omega]^1$			$-\psi[\mu]^v[\omega]$		
		$1 \times$	$+h \times$	$+h^2 \times$	$1 \times$	$+h \times$	$+h^2 \times$
1	•	0	0	$\frac{1}{12}\omega^2$	0	0	$\frac{1}{12}\omega^2$
2	⋮	0	0	$\frac{1}{12}\omega^2$	0	0	$\frac{1}{9}\omega^2 - \frac{1}{36}\mu^2$
	⋮	0	0	$\frac{1}{12}\omega^2$	0	0	$\frac{1}{12}\omega^2 + \frac{1}{108}\mu^2$
3	∇, ∇, ∇	$-\frac{1}{2}$	0		$-\frac{1}{2}$	0	
	⋮, ⋮	$-\frac{1}{2}$	0		$-\frac{1}{2}$	0	
4	∇, ∇, ∇, ...	-1			$-\frac{10}{9}$		
	∇, ∇, ∇, ...	-1			$-\frac{10}{9}$		
	∇, ∇, ∇, ...	-1			-1		
	⋮, ⋮, ⋮	-1			-1		

The overall error of TR - Radaul

$$\begin{aligned}
 b_4 \left(\phi[\omega]^1 + \psi[\mu]^v[\omega] \right) &= \frac{1}{72} \left[-\omega^2 F(\ddagger) \right. \\
 &+ \left(F(\ddagger) - \frac{1}{3} F(\ddagger) \right) \mu^2 + \frac{1}{3} F(\nabla \nabla) + F(\nabla \nabla^*) \\
 &\left. + F(\nabla \nabla^*) + \frac{1}{3} F(\nabla \nabla^*) + F(\nabla \nabla) + F(\nabla \nabla^*) + F(\nabla \nabla) + F(\nabla \nabla^*) \right]
 \end{aligned}$$

The overall error of TR - LobattallIA

Leading residual term

$$\begin{aligned}
 b_5 \left(\phi[\omega] \mathbf{1} + \psi[\mu] \nu[\omega] \right) &= -\frac{1}{72} \left[F(\nabla \bullet) + F(\nabla \bullet) + \frac{1}{2} F(\ddot{\bullet}) \right] \omega^2 \\
 &+ \frac{1}{720} \left[\frac{1}{4} F(\nabla \bullet) + \frac{1}{2} F(\nabla \bullet) + \frac{1}{4} F(\nabla \star) - F(\ddot{\bullet}) - F(\ddot{\bullet}) \right] \mu^2 \\
 &- \frac{1}{720} \left[\frac{1}{4} F(\nabla \nabla \bullet) + F(\nabla \nabla \star) + \frac{3}{2} F(\nabla \nabla \star) \right. \\
 &\left. + F(\nabla \nabla \star) + \frac{1}{4} F(\nabla \nabla \star) + \frac{3}{2} F(\ddot{\nabla} \bullet) + 3F(\ddot{\nabla} \star) + \dots \right]
 \end{aligned}$$



Analysis of the error

Three approaches

- scheme-level approach
- method-level approach
- a priori knowledge

1. The scheme-level approach

Goal : annihilate (or minimize) the leading term of the error of the entire DC scheme.

- one first applies a classical method (such as trapezoidal rule)
- with this numerical solution locally appropriate values for the parameters ω and μ are determined.
 - problems with 2 components at most : **annihilate** the leading error term
 - problems with more than 2 components : **minimize** the leading error term
- Once these values have been obtained, the entire EFDC scheme is applied, giving rise to a method of order
 - $p + r + 1$ for problems with 2 components at most
 - $p + r$ for problems with more than 2 components



1. The scheme-level approach

This algorithm has following properties:

- extra work involved to obtain an initial solution (with a classical method) to estimate values for the parameters ω_k and μ_k .
- computation of the error expression is rather expensive (the higher the order, the higher the number of elementary differentials involved)
- The values for ω_k and μ_k that are obtained in this way, depend upon the specific methods used and **they do not necessarily reflect the nature of the problem.**
- The raise of order is only possible iff $r < p$. For $r = p$, it can be shown that there is an uncontrollable error accumulation (such that the order remains $p + r$).




1. The scheme-level approach

$$\epsilon y'' = y \quad y(0) = 1 \quad y(1) = 0 \quad \epsilon = 0.3$$

$$y(x) = \frac{e^{\frac{x}{\sqrt{\epsilon}}} - e^{\frac{2-x}{\sqrt{\epsilon}}}}{1 - e^{\frac{2}{\sqrt{\epsilon}}}} \quad (\text{test set of J. Cash, Problem 1})$$

This problem can be solved **exactly** by any EF method if the parameter is chosen as $1/\sqrt{\epsilon} \approx 1.8$, but in the scheme-level approach $(\omega_k, \mu_k) \approx (1, -0.7 i)$



2. The method-level approach

Goal : annihilate (or minimize) the leading term of the error of the base method and of the error estimator separately

- one first applies a classical method (such as trapezoidal rule)
- with this numerical solution locally appropriate values for the parameter ω (of the base method) are determined and the EF base method is applied
- with this numerical solution locally appropriate values for the parameter μ (of the error estimator) are determined
- the second step of the DC scheme is applied with the EF base method and the EF error estimator

This gives rise to a method of order

- $p + r + 1$ for scalar problems
- $p + r$ for non-scalar problems



2. Method level approach

This algorithm has following properties:


- still some extra work involved to obtain estimate values for the parameters ω_k and μ_k .
- the values for ω_k and μ_k that are obtained in this way, do reflect the nature of the problem.
- The raise of order is also possible if $r = p$.

2. Method level approach

A Bernoulli-problem

$$y' = \frac{2y + xy^4}{6} \quad y(0) = -2$$

$$y(x) = \frac{-2}{(4x - 4 + 5e^{-x})^{\frac{1}{3}}}$$






2. Method level approach

$$\epsilon y'' = y \quad y(0) = 1 \quad y(1) = 0 \quad \epsilon = 0.3$$

$$y(x) = \frac{e^{\frac{x}{\sqrt{\epsilon}}} - e^{\frac{2-x}{\sqrt{\epsilon}}}}{1 - e^{\frac{2}{\sqrt{\epsilon}}}} \quad (\text{test set of J. Cash, Problem 1})$$

This problem can be solved exactly by any EF method if the parameter is chosen as $1/\sqrt{\epsilon} \approx 1.8$. In the method-level approach $(\omega_k, \mu_k) \approx (1.8, 1.8)$



2. The method level approach

$$y'' = -y - y^3 + 0.002 \cos(1.01x) \quad (\text{Duffing problem})$$

$$y(0) = 0.200426728067 \quad y(2) = -0.08668702310$$

Approximate solution:

$$\begin{aligned} y(x) = & 0.200179477536 \cos(1.01x) + 0.000246946143 \cos(3.03x) \\ & + 0.000000304014 \cos(5.05x) + 0.000000000374 \cos(7.07x) \\ & + \dots \end{aligned}$$

EF methods can produce very accurate results for this problem if the parameter is approximately equal to 1.01


2. The method level approach

$$y'' = -y - y^3 + 0.002 \cos(1.01x) \quad (\text{Duffing problem})$$

$$y(0) = 0.200426728067 \quad y(2) = -0.08668702310$$

EF methods can produce very accurate results for this problem if the parameter is approximately equal to 1.01.

The computed values for ω_k and μ_k vary in the integration interval from 1.5 to 1





3. A priori knowledge

Suppose we already have a good estimate for the EF parameter ω .


- One then immediately applies an EF method (such as EF trapezoidal rule) with that estimate
- with this numerical solution locally appropriate values for the parameter μ (of the error estimator) are determined
- the second step of the DC scheme is applied with the original EF base method and the EF error estimator



3. A priori knowledge

$$y'' = - \left(100 + \frac{1}{4x^2} \right) y \quad y(1) = J_0(10), \quad y(2) = \sqrt{2}J_0(20)$$

$$y(x) = \sqrt{x}J_0(10x) \quad (\text{Bessel problem})$$



DC : classical DC scheme

Algorithm 1 : a priori fixed ω -values : $\omega_k = 10^i$

Algorithm 2 : the method-level approach



The end

Thank you for your attention